



A study to establish the reference values for two minute walk distance in school going children

Nawaj Pathan

Assistant Professor, MGM School of Physiotherapy, Aurangabad, Maharashtra, India

Abstract

Background: The idea to use functional scales to evaluate the prognosis of children is widely being facilitate by the International Classification of Functioning, Disabilities & Health, hence the present study was to examine responsiveness, of 2 Minute Walk Test (MWT) in school going children.

Purpose of the Study: This current study aims to establish the reference values for 2MWT in healthy school going children.

Objective of study: To establish reference values for 2 Min Walk distance in healthy Indian children of 7 to 10 years age.

Method & material: A total of 82 school going children participated in the study aged between 7 to 10 years. Out of which 41 were normal weight and 41 were obese students. They were divided equally into two groups based on their BMI. Group A comprised of obese and group B comprised of normal weight participants. Two-minute walk test was performed by the participants on a obstacle free 15 m distance. Distance covered by the participants was measured and documented after the test.

Results: The mean 2MWD was 158.268 ± 16.0209 meters in obese children whereas in normal weight children it was 185.2195 ± 13.81771 . 2MWD had shown excellent correlation with weight. $P=0.0001$ is considered as significant.

Conclusion: The average distance covered by healthy children is approximately 185.2195 ± 13.81771 meters in two minute which is less in obese group 158.268 ± 16.0209 meters.

Keywords: 2 minute walk test, functional mobility, obesity, school going children

1. Introduction

Childhood obesity is a prominent & critical public health concern ^[1] The percentage of overweight and obesity problems in school going boys and girls increasing. This rising percentage of overweight reached 15.2% ^[2] Walking is an important and fundamental task, characterized by a complex interrelated task between neuromuscular, environmental factors. Overweight and obese children have a typical gait pattern which is short ^[3] insufficient postural control ^[4] the elevated impact of force on lower limbs ^[5], malalignments of lower extremities ^[6] which may interfere in their important day to day life activities i.e. Walking ^[7].

Low levels of physical activity associated with both physiological and biomechanical factors, which are mainly responsible for changes in gait patterns in obese and overweight children, have been observed ^[8]. Such a distinction not only offers quantification between obese and overweight, but also offers an important details on the variables in determining their mobility ability and physical activities. Several musculoskeletal conditions occur as a result of being overweight which greatly reduced mobility in the paediatric population ^[9] Gait pattern in obese children changed mainly in their temporospatial components mainly walking speed, stride time and length, step length timing, duration of the stance phase, and swing phase also typically slowed down ^[10] Obese children have slower walking speed, shorter strides, a large period of the support phase in the gait cycle, low stability, and narrow or reduced joint angles. Altered walking velocity prolongs the stance phase. Obese children associated with flat feet fatigued more easily than normal weighted children ^[11] Dynamic gait alterations in these children increase compressive forces in the medial compartment of the knee which is considered as

pathological.⁹ There are a lot of walk tests available which measures the distance covered in time by the individual, but most of the walk tests have drawbacks such as participants who cannot complete the tests till the end and fewer rest intervals,¹² so that is why the need of 2 min walk test has been generated for various age groups. 2 min walk test has been widely used in stroke, COPD, lower limb amputees to measure the endurance level.

In their study, Kieseier and Pozzilli suggested that 2 MWT assess walking fatigue and distance limitations and also found that compared to 6 MWT, 2 MWT is a more convenient measure of uninterrupted walking or usual walking performance.¹³ As per our knowledge & literature search there are few studies available in the Indian paediatric population to measure walking speed, distance limitations in overweight, obese and normal-weight children, hence this study has been taken up to investigate the usefulness of 2MWT to measure walking speed in these children.

Methodology

Study Design

Cross Sectional Study.

Sample Size

82

Inclusion Criteria

Being able to walk 15 m without a walking aid. Both genders were included Age group ranges between 7-10 No orthopaedic, surgical, or neurological problems affecting the gait. BMI ranges between 18 and 25 kg/m² according to the WHO's classification & guidelines.

Exclusion Criteria

Children having foot deformities such as ulcers, recent injuries. Impaired sensations in lower limbs. Had history of vascular problems which affecting their gait abilities. Any systemic illness affecting their waking abilities.

Instruments



Fig 1: Stopwatch



Fig 2: Inch tape



Fig 3: Weighing machine



Fig 4: Two Cones

Procedure

Schools in Aurangabad were selected through simple randomization. Prior to the data collection written consent was taken from the participant’s parents. After getting the consent, students were briefly demonstrated about the test. Demographic data like name, age, gender, height and weight of all the participants was documented before performing the test. Each student’s BMI was calculated using the formula weight in kg divided by height in m². After obtaining the BMI participants were divided into two groups. Group A comprised of obese and Group B comprised of normal students. Each group had 41 participants in it. An obstacle free distance of 15m was used for the test. Prior to test instructions were given to the participants to walk at normal speed and pace and not to run or jump. Two trial tests were performed by the students and the best of two was considered. Individual student walked for 2 minutes, after finishing the test distance covered by the participant in 2 minutes was documented.

Outcome measures
Body mass index ^[17]

$$BMI = \text{Weight kgs} / \text{Height m}^2$$

Table 1

Weight Status Category	Percentile Range
Underweight	Less Than 5 th Percentile
Normal Or Healthy Weight	5 th Percentile To Less Than 85 th Percentile
Overweight	85 th To Less Than 95 th Percentile
Obese	Equal To Or Greater Than The 95 th Percentile

2 min walk test ^[18]

Two Minute Walk Test

Name of Partici

Date

Distance ambulated in 2 minutes:-

Results

Table 1: Comparison of Mean Age in Groups

Age	Obese	Normal	PERCENTAGE	
			Obese %	Normal %
7	8	6	19.51%	14.63%
8	15	10	36.58%	24.39%
9	11	18	26.82%	43.90%
10	7	7	17.07%	17.07%
Total	41	41		
(Mean ± SD)	158.26 ± 16.02	185.21 ± 13.81		




Fig 5: Comparison of Mean Age in Groups

Table 2: Comparison of Gender in Groups

Gender			Percentage	
	Obese	Normal	Obese %	Normal %
Male	15	22	36.58	53.65
Female	26	19	63.41	46.34
Total	41	41		




Fig 6: Comparison of Gender in Groups

Table 3: Comparison of mean distance in groups

	Obese (mean ± sd)	Normal (mean ± sd)	T value	P value
Distance	158.268 ± 16.0209	185.2195 ± 13.81771	7.96	<0.0001

Discussion

This era of technology where the deskbound lifestyle such as surplus use of Mobiles, computer, video gaming and other electronic gadgets along with the dietary changes such as excess consumption of high-energy, high fat diets mainly fast food, snacks, soft drinks. Deterioration in dietary quality associated with inactive behaviour plays a crucial role in the mounting obesity in children, which further precedes to risks of non-communicable diseases like Diabetes, cardiovascular conditions, insulin resistance, low-grade inflammation, increased risk of cardio-metabolic abnormalities [19] Cancer, Depression, Anxiety [20] Asthma along with various other Musculoskeletal issues etc [21] The demand for clinical assessment to bench mark the functional capacity of such overweight or obese children is increasing. The 6-minute walk test (6MWT) is a straightforward, practical, reliable and valid test for evaluating sub maximal exercise capacity in healthy kids but it requires more time, training and instrumentation to perform. Children even drop out before they could complete 6 minutes as they are fatigued or due to Cardiovascular and musculoskeletal issues that often arise during the performance of this test. Therefore 6MWT might not function as an efficient measure for observance of practical ability and illness progression in the children. However, the 2-Minute walk Test (2MWT) is of shorter duration, objective, easy to administer in paediatric populations, [22] which provides a uniformity in analysis of functional capacities in paediatric participants and it highly co-relates with 6MWT [23] The 2MWT could effectively replace the 6MWT in this population and predict

the same as there is an increase in prevalence of adiposity between the ages 7-11 years^[24]. The mid-late childhood which is around age group between 7-11 years could get benefit and larger attention in future obesity bar interventions^[25, 26]. So To evaluate the functional mobility the total distance covered by the children's aging in this age group was measured and evaluated with the assistance of two min walk test. Students were categorized and selected as normal weighing (NW) or overweight and obese (OW) on the basis of BMI as per the guidelines by CDC^[17]. Overweight is defined clinically as body mass index (BMI) \geq 85th percentile and $<$ 95th percentile with obesity classified as \geq 95th percentile. Students were explained and demonstrated 2 min walk test properly. To avoid any mishap safety precautions were taken. The walkways employed in our study measured 15m with 2 cones placed on the both the ends as a turning point. Due to a variety of constraints involved during the testing in school setting, the test was performed with every student on an individual basis to avoid students matching pace with their peers. Each student was allotted a physiotherapist to document and evaluate individually thus competition among students were eliminated and findings taken into considerations. Two trials were performed by each student with a time frame of 20 second gap in between two trials, the best of two was selected. The Normal Weight and Obese children's were segregated with 41 students in each group, in group A there were 15 males and 26 females and in group B there were 22 males and 19 females as participants. The mean distance covered by the Obese children i.e. group A was 158.268 ± 16.0209 while the Normal Weight i.e. group B covered mean distance of 185.2195 ± 13.81771 with p value <0.0001 thus indicating significance. In this trail, it was observed that Obese children walked less distance with a speed that decreased over a period of time as compared to Normal Weight children. We have also found the prevalence rate of obesity in girl was more as compared to boys. Results of this research suggested a direct correlation between weight status and functional capacity. The walking speed might be affected resulting from altered biomechanics, kinematic and gait parameters that were altered by changes in anthropometric measures^[28] or due to decrease in cardiovascular capacity thereby minimizing the efficiency to perform but the root cause couldn't be specified. Roya Kelishadi et. Al. reported that BMI and physical activity are co-related. It was found that children's with insufficient physical activity had higher BMI and weight compared to more physically active children's. The government should introduce a national strategy to encourage physical activity, as the classroom is the location where children spend most of their time. Addition to this, parents can encourage more physical activity by reducing the screen time and by making physical activity more helpful and enjoyable^[30].

Conclusion

The present study has concludes that the 2MWT is a useful tool to measure the walking speed in obese and normal weight children.

Acknowledgement

All the authors wants to thank all children participated in the study, School Principal Mrs. Sabahat Khan madam who permitted us to conduct this study.

Conflict Of Interest

None

Source of Funding

None.

References

1. Karnik S, Kanekar A. Childhood Obesity: A Global Public Health Crisis. *Int. J. Prev. Med.* 2012; 3:1-7.
2. Xiyan Zhang, Fengyun Zhang. Prevalence of Overweight and Obesity among Primary School-Aged Children in Jiangsu Province, China, 2014-2017. *Overweight/Obesity in Jiangsu Province, 2018, 13(8).* (Accessed 5/3/2020)
3. Lerner ZF, Board WJ, Browning RC. Effects of Obesity on Lower Extremity Muscle Function During Walking At Two Speeds. *Gait Posture.* 2014; 39:978-984.
4. Deforche BI, Hills AP, Worringham CJ, Davies PS, Murphy AJ, Bouckaert JJ, et al. Balance And Postural Skills In Normal-Weight And Overweight Prepubertal Boys. *Int. J. Pediatr. Obes.* 2009; 4:175-182.
5. Tsiros MD, Coates AM, Howe PR, Grimshaw PN, Walkley J, Shield A, et al.. Knee Extensor Strength Differences In Obese And Healthy-Weight 10-To 13-Year-Olds. *Eur. J. Appl. Physiol.* 2013; 113:1415-1422.
6. Shultz SP, D'Hondt E, Lenoir M, Fink PW, Hills AP. The Role of Excess Mass in the Adaptation Of Children's Gait. *Hum. Mov. Sci.* 2014; 36:12-19.
7. Andrew P Hills, Anthony W Parker, Gait Characteristics In Obese Children, *Arch Phy. Medi. Rehabilitation*, 1991.
8. Sarah P, Shultz Raymondc Browning. Childhood Obesity and Walking: Guidelines And Challenges. *Internatinal Journal Of Pediatric Obesity.* 2011; 6:5-6. [Http://Www.Tandfonline. Com/Doi/Abs/10.3109/17477166.2011.590202](http://Www.Tandfonline.Com/Doi/Abs/10.3109/17477166.2011.590202) (Accessed 6/12/19).
9. S.P. Shultz, J. Anner And A. P.Hills. Paediatric Obesity, Physical Activity and The Musculoskeletal System. *Obesity*, 2009, 10. [Http://Www.Ncbi.Nlm.Nih.Gov/M/Pubmed/194660114 /](http://Www.Ncbi.Nlm.Nih.Gov/M/Pubmed/194660114/) (Accessed 6/12/19).
10. Wall JC. Charteris J. Turnbull GI. Two Steps Equals One Stride Equals What? The Applicability Of Normal Gait Nomenclature To Abnormal Walking Patterns. *Clin. Biomech.* 1987; 2:119-125.
11. Songhua Yan, Weiyanrenetal. Gait Characteristics Of Overweight And Obese Children With Different Ages. *IEEE*, 2010, 25(5). (Accessed 5/3/2020).
12. Mchardy GJR, Gupta SP, Mchardy GJR: Twelve-Minute Walk Test For Assessing Disability In Chronic Bronchitis. *Br Med J.* 1976; 1:822-823.
13. David A, Scalzitti Etal. Validation of The 2-Minute Walk Test With The 6-Minute Walk Test And Other Functional Measures In Persons With Multiple Sclerosis. *International Journal Of MS Care*, 2018. (Accessed 5/3/2020).
14. Nuttall Frank Q. "Body Mass Index: Obesity, BMI, And Health: A Critical Review." *Nutrition Today.* 2017; 50(3):117-128. Doi: 10.1097/ NT.000 000000000092.
15. Bohannon Richard W. Normative Reference Values For The Two-Minute Walk Test Derived By Meta-Analysis. *Journal Of Physical Therapy Science.* 2017;

- 29(12):2224-2227. Doi:10.1589/ Jpts.29.2224.
16. Harrington DM, Staiano AE, Broyles ST, Gupta AK, Katzmarzyk PT. BMI Percentiles For The Identification Of Abdominal Obesity And Metabolic Risk In Children And Adolescents: Evidence In Support Of The CDC 95th Percentile. *Eur J Clin Nutr.* 2013; 67(2):218-222. Doi:10.1038/Ejcn.2012.203
 17. Pin TW, Choi HL. Reliability, Validity, And Norms Of The 2-Min Walk Test In Children With And Without Neuromuscular Disorders Aged 6-12. *Disabil Rehabil.* 2018; 40(11):1266-1272. Doi:10.1080/09638288.2017.1294208
 18. Gulati Achal, Hochdorn Alexander, Paramesh, Haralappa, Paramesh Elizabeth, Chiffi Daniele, *et al.* Physical Activity Patterns Among School Children In India. *Indian Journal Of Pediatrics.* 2014; 81:10-1007/S12098-014-1472-X.
 19. Krul M, Van Der Wouden JC, Schellevis FG, Van Suijlekom-Smit LW, Koes BW. Musculoskeletal Problems In Overweight And Obese Children. *Ann Fam Med.* 2009; 7(4):352-356. Doi:10.1370/Afm.1005
 20. Xu Shumei, Ying Xue. "Pediatric Obesity: Causes, Symptoms, Prevention And Treatment." *Experimental And Therapeutic Medicine.* 2016; 11(1):5-20. Doi:10.3892/ Etm.2015.285
 21. Witherspoon JW, Vasavada R, Logaraj RH. Two-Minute Versus 6-Minute Walk Distances During 6-Minute Walk Test In Neuromuscular Disease: Is The 2-Minute Walk Test An Effective Alternative To A 6-Minute Walk Test?. *Eur J Paediatr Neurol.* 2019; 23(1):165-170. Doi:10.1016/J.Ejpn.2018.10.001
 22. Lauren Reid MD, Penny Thomson Bsc, Markus Besemann MD, Nancy Dudek MD. *Med Going Places: Does The Two-Minute Walk Test Predict The Six Minute Walk Test In Lower Extremity Amputees? (N.D).* *J Rehabilmed,* 2015, 256-261
 23. Bohannon RW, Bubela D, Magasi S. Comparison Of Walking Performance Over The First 2 Minutes And The Full 6 Minutes Of The Six-Minute Walk Test. *BMC Res Notes.* 2014; 7:269. Published 2014 Apr 25. Doi:10.1186/1756-0500-7-269
 24. Hughes AR, Sherriff A, Lawlor DA, Ness AR, Reilly JJ. Incidence of Obesity During Childhood And Adolescence In A Large Contemporary Cohort. *Prev Med.* 2011; 52(5):300-304. Doi:10.1016/J.Ypmed.2011.02.014
 25. Von Kries R, Reulen H, Bayer O, Riedel C, Diethelm K, Buyken AE. Increase In Prevalence Of Adiposity Between The Ages Of 7 And 11 Years Reflects Lower Remission Rates During This Period. *Pediatr Obes.* 2013; 8(1):13-20. Doi:10.1111/J.2047-6310.2012.00084.X
 26. Horsak B, Schwab C, Baca A. Effects Of A Lower Extremity Exercise Program On Gait Biomechanics And Clinical Outcomes In Children And Adolescents With Obesity: A Randomized Controlled Trial. *Gait Posture.* 2019; 70:122-129. Doi:10.1016/J.Gaitpost.2019.02.032
 27. Himes JH. Challenges of Accurately Measuring And Using BMI And Other Indicators Of Obesity In Children. *Pediatrics.* 2009 124(1):S3-S22. Doi:10.1542/Peds.2008-3586D
 28. Kelishadi R, Qorbani M, Djalalinia S. Physical Inactivity And Associated Factors In Iranian Children And Adolescents: The Weight Disorders Survey Of The CASPIAN-IV Study. *J Cardiovasc Thorac Res.* 2017; 9(1):41-48. Doi:10.15171/Jcvtr.2017.06
 29. Colberg SR, Sigal RJ, Fernhall B. Exercise And Type 2 Diabetes: The American College Of Sports Medicine And The American Diabetes Association: Joint Position Statement. *Diabetes Care.* 2010; 33(12):E147-E167. Doi:10.2337/ Dc10-9990
 30. Weiler Richard, Allardyce Sam, Whyte Greg, Stamatakis Emmanuel. Is The Lack Of Physical Activity Strategy For Children Complicit Mass Child Neglect?. *British Journal Of Sports Medicine.* 2013; 48:10-1136/Bjsports-2013-093018.S