

Studies on the antimicrobial effect of various plant extracts against Human pathogens and search for the New Drugs

^{1, 2}Kumaran T, ² Citarasu T

¹ Department of Zoology, Muslim Arts College, Thiruvithancode, Kanyakumari District 629174, India.

² Centre for Marine Science and Technology, Manonmaniam Sundaranar University, Rajakkamangalam 629502, India.

Abstract

Medicinal plants are considered as an important source of potentially useful structures for the development of new chemotherapeutic agents. In the present study, three selected Indian medicinal plants such as *Hybanthus enneaspermus*, *Asparagus racemosus* and *Curcuma longa* were analyzed for phytochemical constituents and tested for antibacterial activity against pathogens. In most of the samples all the phytochemicals i.e. reducing sugar, Terpenoids, Flavonoids, Saponins, alkaloids, cardiac glycosides, carbohydrates and phytosterols were present. Methanol and aqueous extracts of the samples of all plants were evaluated for *in vitro* antibacterial activity against *Escherichia coli*, *Staphylococcus aureus* and *Pseudomonas aeruginosa* by agar plate well diffusion method. All the plant extracts showed significant antibacterial activity against the tested organisms. However, methanol extracts showed maximum inhibitory effect from this *Hybanthus enneaspermus* and *Asparagus racemosus* were showed maximum bacterial inhibition than other plants.

Keywords: Medicinal plants, Phytochemical, antibacterial activity, pathogens

Introduction

Medicinal and aromatic plants constitute a major source of natural organic compounds and serve as the immediate source of drugs (Dohbal, 1994). The medicinal plants generally contain number of compounds that may be potential natural antimicrobial agents which may serve as alternative, effective, cheaper and safe antimicrobial agents for the treatment of common microbial infections (Schimmer *et al.*, 1994).

Antibiotics such as ampicillin, bacitracin, lincomycin, chlortetracycline and streptomycin promote growth because of an effect on the microflora in the gastrointestinal tract (Coates *et al.*, 1963; deMan, 1975). The phasing out of antibiotic growth promoters will affect the animal industry at large. To minimize the loss in growth, there is a need to find alternatives to antibiotic growth promoters. There are a number of non-therapeutic alternatives to antibiotic growth promoters, including enzymes, (in) organic acids, probiotics, herbs, immunostimulants and specific management practices (McEwen and Fedorka-cray, 2002).

More recently, medicinal plant extracts were developed and proposed for use in food as natural antimicrobials (Hsieh *et al.*, 2001). The natural plant products have been reported to promote various activities like antistress, growth promoting, appetizing, tonic, immunostimulation, aphrodisiac and antimicrobials in the finfish and shrimp larviculture due to the active principle natures such as alkaloids, flavanoids, pigments, phenolics, terpenoids, steroids and essential oils. (Citarasu *et al.*, 1998; Citarasu *et al.*, 2002). Thus, in addition to enhancing the immune response, the antimicrobial effects of these herbs are beneficial to the host (Citarasu, 2012). The present study was conducted to determine the effect of different medicinal plant (herbs) extracts as a possible alternative to antibiotics.

Asparagus racemosus (Shatavari) has been described as a rasayana herb and has been used extensively as an adaptogen to increase the non-specific resistance of organisms against a variety of stresses. Besides use in the treatment of diarrhoea and dysentery, the plant also has antioxidant, immunostimulant, anti-dyspepsia and antitussive effects (Bopana and Saxena 2007).

Hybanthus enneaspermus is used to determine antibacterial activity against some urinary tract pathogens. This plant has various biological activities such as anti-inflammatory, antifussive, antiplasmodial (Singh *et al.*, 1980).

Curcuma longa (Turmeric), a perennial herb, has along tradition of use in the Chinese and Ayurvedic systems of medicine. Curcuminoids, a group of phenolic compounds isolated from the roots of *C. longa*, exhibited a variety of beneficial effects on health and has the ability to prevent certain diseases (Joe *et al.*, 2004). In South Asia, the rhizomes from *C. longa*, are considered to have natural medicinal properties, including antibacterial, anti-inflammatory.

Material and Methods

Selection of medicinal plants for this study

Three medicinal plants including *Hybanthus enneaspermus*, *Asparagus racemosus* root (Shatavari), *Curcuma longa* rhizomes (Turmeric), were utilized in this study. These plants have previously been reported to have antibacterial activity against different bacterial strains.

Preparation of Extracts

Processing of the selected plant materials

Collected plant materials were shade dried with in temperature range of 28-35 °C. The drying process was continued to reduce moisture less than 14%. After drying, the plant materials were minced with wooden knife feeding into a

grinder, minced materials were made into powder using teeth mills and sieved, and then the powder was stored in airtight container and kept at room temperature until further use.

Extraction of selected plant material powder

10g of dried powder was taken and boil at 100°C for 2 hours, then filter the extracts, supernatant was collected and residues were discarded. The collected supernatant was condensed in water bath, and condensate was extracted again with methanol. The methanolic extract was concentrated in rotary evaporated under reduced pressure at room temperature 45-50 °C, in order to avoid evaporation of plant metabolism. Aqueous extract was concentrated and stored at 4 °C.

In vitro antibacterial studies

Antimicrobial susceptibility studies

Inhibition of microbial growth was tested by using the paper disc agar diffusion method (Bauer-Kirby Method) while the MIC was determined by the dilution (both micro and macro) method (de Paiva *et al.*, 2003). Standard aseptic microbiological methods were followed throughout this antibacterial study. Microorganisms such as *E. coli*, *P. aeruginosa* and *S. aureus* (Centre for Marine Science and Technology, Manonmaniam Sundaranar University, Rajakkamangalam), were obtained.

Screening by agar well diffusion method

The fractions eluted from the column purification were screened by agar disc diffusion method (Bauer and Kirby, 1966). Disc of filter paper (5 MM) impregnated with herbal extracts were placed on an agar plate that was heavily and uniformly inoculated (lawn) with an actively growing culture of the organisms. The medium of choice was Muller Hinton Agar for *P. aeruginosa*, *S. aureus* and *E. coli*. The dynamics and timing of antimicrobial agent diffusion to establish a concentration gradient coupled with the growth of organisms over 16 – 24 hour duration is critical for reliable results.

Minimum inhibitory concentration (MIC) and Minimal Bacterial Concentration (MBC)

Of the 4 plants tested, only those that showed antibacterial activity against some of the selected pathogens were selected for further tests to calculate their MIC by dilution method. The Minimum bacterial concentration (MBC) (Jacobs and Demott, 1994) was determined by plating the culture from the tube without the growth of microorganisms. To determine the minimum bacterial concentration, aliquots of one loop full of the two lowest concentrations which inhibited bacterial growth, were streaked on petriplates containing Muller Hinton agar. After as overnight incubation at 37 °C the plates were evaluated by comparing them with control plates containing bacteria without test compounds. The lowest concentration that gave no visible growth was taken as MBC.

Results

Antibacterial screening

Many of the plant species have been documented pharmacologically and clinically which are endowed in phytochemistry with marked activity on human pathogenic bacteria. The methanol extract of the medicinal plants *H. enneaspermus*, *A. racemosus* root (Shatavari) *C. longa* rhizomes (Turmeric) showed high level antibacterial activity

against *P. aeruginosa* and *S. aureus* at specific doses mentioned.

Antibacterial screening for Agar well and Disc diffusion method

The antibacterial activities of the selected extracts (Zone of inhibition of mm diameter) were given in the Table 1. Among the different active extracts, *A. racemosus* controlled the pathogen of 12.30, 12 and 14 mm of zone of inhibition in *P. aeruginosa*, *S. aureus* and *E. coli* respectively. The *H. enneaspermus* extract controlled the range of 9 to 13 mm of zone of inhibition in *P. aeruginosa*, *S. aureus* and *E. coli* respectively. The methanol extract of the medicinal plants *C. longa*, showed no antibacterial activity against *P. aeruginosa*, *S. aureus* and *E. coli* at specific doses mentioned in Table 1 zone of inhibition respectively.

Table 1: Antibacterial screening of the active fractions of different herbals extracts against pathogens

Sl. No	Pathogens	Zone of inhibition (mm) of different active fraction of the herbal extracts		
		<i>H. enneaspermus</i>	<i>A. racemosus</i>	<i>C. longa</i>
1	<i>P. aeruginosa</i>	12.30 ± 0.94	9.30 ± 1.04	3.12 ± 1.02
2	<i>S. aureus</i>	12.0 ± 0.55	10.33 ± 1.63	1.26 ± 1.16
3	<i>E. coli</i>	14.0 ± 0.95	12.4 ± 0.63	2.08 ± 0.14

Minimum Inhibitory Concentration (MIC) and Minimal Bacterial Concentration (MBC)

The growth rate of the selected bacterial isolates at different hours was given in the Fig 1. The MIC and MBC of the above three extracts against the pathogens were given in the Tables 2a to 2c. The MICs are 60, 70 and 50 µg in the three extracts against *P. aeruginosa*. For *S. aureus*, the MIC is 60 to 70 µg of the selected plant extracts. The MIC is further decreased to 40 to 60 µg against the *E. coli* of the selected extracts. The least minimum inhibitory concentration was detected against the *E. coli*.




Fig 1: Growth rate of the selected bacterial isolates at different hours

Table 2a: Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal concentration (MBC) of the active extract against the pathogen *Pseudomonas aeruginosa*

Con (µg)	Active Herbal Extract fractions					
	<i>H. Enneaspermus</i>		<i>A. racemosus</i>		<i>C. longa</i>	
	MIC	MBC	MIC	MIC	MIC	MBC
10		++++		++++		++
20		++++		++++		+
30		++		++		-
40		++		++		-
50		++		++	10	-
60	60	+		+		-
70		-	70	+		-
80		-		-		-

- : No Growth; +: Minimum Growth; ++: Intermediate Growth; >+: Maximum Growth

Table 2b: Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) of the active extract against the pathogen *Staphylococcus aureus*

Con (µg)	Active Herbal Extract fractions					
	<i>H. enneaspermus</i>		<i>A. racemosus</i>		<i>C. longa</i>	
	MIC	MBC	MIC	MIC	MIC	MBC
10		++++		++++		+
20		++		+++		+
30		++		++		-
40		++	60	++		-
50		++		++	10	-
60		-		-		-
70	70	-		-		-
80		-		-		-

- : No Growth; +: Minimum Growth; ++: Intermediate Growth; >+: Maximum Growth

Table 2c: Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) of the active extract against the pathogen *E. coli*.

Con (µg)	Active Herbal Extract fractions					
	<i>H. enneaspermus</i>		<i>A. racemosus</i>		<i>C. longa</i>	
	MIC	MBC	MIC	MIC	MIC	MBC
10		++++		++++		+
20		++		++++		-
30		++		++		-
40	40	+		++		-
50		-		-	10	-
60		-	60	-		-
70		-		-		-

- : No Growth; +: Minimum Growth; ++: Intermediate Growth; >+: Maximum Growth

Discussion

The antimicrobial effect of the medicinal plants is well documented (Valero and Salmeron, 2003). The results of different studies provide evidence that some medicinal plants might indeed be potential sources of new antibacterial agents even against some antibiotic-resistant strains (Kone *et al.*, 2004). In this study, using the disk diffusion method it was observed that extracts of Shatavari and thyme produce antibacterial activity against both gram negative and gram positive pathogens. The aqueous extract of *A. racemosus* exhibited moderate level antimicrobial activity against *E.coli*, *P. aeruginosa* and *V. parahaemolyticus* (Kumaran and Citarasu 2015).

The antimicrobial activities of the Shatavari plant extracts against the bacterial strains examined were assessed by the presence or absence of inhibition zones. The Shatavari extract was found to be effective against *E. coli*, *P. aeruginosa*, and *S. aureus*. This effect is in agreement with other researchers regarding the antibacterial effect against *E. coli* however there is a difference in the concentration of extract of Shatavari at which we found antibacterial activity (Rajpal 2002). Using the disk diffusion method, the concentration at which antibacterial activity was found was much higher than that of mentioned by the above mentioned authors. The MIC results of Shatavari by dilution method in our studies support an earlier finding by Rajpal 2002. In our studies the *H. enneaspermus* extract showed antibacterial activity. This result supports the findings of many authors (Nevas *et al.*, 2004). *H. enneaspermus* was found to be effective against *E. coli* and *P. aeruginosa* but not *S. aureus* by the disk diffusion method.

There was no antibacterial activity in extracts of *C. longa* against the tested pathogens at the specific dose. Our results are contradictory with some researchers who reported antibacterial activity of above plants against gram positive and gram negative bacteria (Kalemba and Kunicka, 2003). This variation may be because of the dose used in this study, the method of extraction of medicinal plants, the method of antibacterial study, the genetic variation of plant, age of the plant or the environment. Our results suggest that gram-positive and gram negative bacteria are generally more sensitive to the spice and herb extracts. This was consistent with the previous studies on other spices and herbs (Ceylan and Fung 2004).

References

- Bauer AW, Kirby WMM, Serris JC, Turck M. Antibiotic susceptibility testing by a standardized single disc method. American Journal of Clinical Pathology. 1966; 45:493-496.
- Bopana N, Saxena S. Asparagus racemosus-ethnopharmacological evaluation and conservation needs. Journal of Ethnopharmacology. 2007; 110(1):1-15,
- Ceylan E, Fung DYC. Antimicrobial activity of spices. J Rap Meth Auto Microbiol. 2004; 12:1-55.
- Citarasu, Sekar TRR, Babu MM, Marian MP. Developing *Artemia* enriched herbal diet for producing quality larvae in *Peaneus monodon*, Asian Fisheries Science. 2002; 15:21-32.
- Citarasu. Natural Antimicrobials in aquaculture, In: Infectious diseases in aquaculture: prevention and control, (Ed. Brain Austin) Woodhead Publishing Ltd, UK, 2012, 419-446.
- Citarasu T, Jayarani TV, Babu MM, Marian MP. Application of biomedicinal products for improving marine shrimp larval production *Aqua-Terr. Annual symposium*, School of Biological Sciences, M.K. University, Madurai. 1998.
- Coates ME, Fuller R, Harrison GF, Lev M, Suffolk SF. A comparison of the growth of chicks in the Gustafson germ-free apparatus and in a conventional environment, with and without dietary supplements of penicillin. Br. J. Nutr. 1963; 17:141-150.

8. De Paiva SR, Figueiredo MR, Aragao TV, Kaplan MA. Antimicrobial activity extracts. *Food Microbiol.* 2003; 18:35-43.
9. De man JC. The probability of most probable numbers. *Eur. J. Appl. Microbiol.* 1975; 1:6778.
10. Dohbal R. Medicinal and aromatic plants. *Curr. Sci.* 1994; 71:397-398.
11. Habsah M, Amran M, Mackeen MM, Lajis NH, Kikuzaki H, Nakatani N, Rahman AA, Ali Ghafar AM. Screening of Zingiberaceae extracts for antimicrobial and antioxidant activities. *J. Ethnopharmacol.* 2000; 72:403-410.
12. Hsieh PC, Mau JL, SH. Antimicrobial effect of various combinations of plant. *In vitro* of plumbagin isolated from Plumbago species. *Mem Inst Oswaldo Cruz.* 2001; 98:959-61.
13. Joe B, Vijaykumar M, Lokesh BR. Biological properties of curcumin-cellular and molecular mechanisms of action. *Crit. Rev. Food Sci. Nutr.* 2004; 44:97-111.
14. Kalemba D, Kunicka A. Antibacterial and antifungal properties of essential oils. *Curr. Med Chem.* 2003; 10:813-829.
15. Kone WM, Atindehou KK, Terreaux C, Hostettmann K, Traore D, Dosso M. Traditional medicine in North Cote-d'Ivoire: screening of 50 medicinal plants for antibacterial activity. *J. Ethnopharmacol.* 2004; 93:43-49.
16. Kumaran T, Citarasu T. Ethnopharmacological investigation and antibacterial evaluation of the methanolic extract of *Asparagus racemosus* (Shatavari). *Tropical Plant Research.* 2015; 2(3):175-179.
17. McEwen SA, Fedorka-Cray PJ. Antimicrobial use and resistance in animals. *Clin. Infect. Dis.* 2002; 34:93-106.
18. Nevas M, Korhonen AR, Lindstrom M, Turkki P, Korkeala H. Antibacterial efficiency of finnish spice essential oils against pathogenic and spoilage bacteria. *J. Food. Prot.* 2004; 67:199-202.
19. Rajpal V. Indian Herbal Pharmacopoeia Standardization of Botanicals, 2002, 2, 2.
20. Schimmer O, Kruger A, Paulin H, Haefele F. An evaluation of 55 commercially plants extract in the Ames Mutagenicity test. *Pharmazie,* 1994; 49:448-451.
21. Singh VP, Sharma SK, Khan VS. Medicinal Plants from Vjain District Madhya Pradesh part II. *Indian Drugs and Pharmaceutical Industry* 1980; 5:7-12.
22. Srinivasan D, Nathan S, Suresh T, Lakshmana Perumalsamy P. Antimicrobial activity of certain Indian medicinal plants used in folkloric medicine. *J. Ethnopharmacol.* 2001; 74:217-220.
23. Valero M, Salmeron MC. Antibacterial activity of 11 essential oils against *Bacillus cereus* in tyndallized carrot broth. *Int. J. Food Microbiol.* 2003; 85:73-81.